

## Clinical Policy: Anakinra (Kineret)

Reference Number: DE.PHAR.244

Effective Date: 01.23

Last Review Date: 01.23

Line of Business: Medicaid

[Coding Implications](#)  
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

Anakinra (Kineret<sup>®</sup>) is an interleukin-1 (IL-1) receptor antagonist.

### FDA Approved Indication(s)

Kineret is indicated for the treatment of:

- Rheumatoid arthritis (RA): Reduction in signs and symptoms and slowing the progression of structural damage in moderately to severely active RA, in patients 18 years of age or older who have failed 1 or more disease modifying antirheumatic drugs (DMARDs). Kineret can be used alone or in combination with DMARDs other than tumor necrosis factor blocking agents.
- Cryopyrin-associated periodic syndromes (CAPS): Treatment of neonatal-onset multisystem inflammatory disease (NOMID).
- Deficiency of interleukin-1 receptor antagonist (DIRA): Treatment of DIRA.

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Kineret is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Rheumatoid Arthritis (must meet all):

1. Diagnosis of RA per American College of Rheumatology (ACR) criteria (*see Appendix F*);
2. Prescribed by or in consultation with a rheumatologist;
3. Age  $\geq$  18 years;
4. Member meets one of the following (a or b):
  - a. Failure of a  $\geq$  3 consecutive month trial of methotrexate (MTX) at up to maximally indicated doses;
  - b. Member has intolerance or contraindication to MTX (*see Appendix D*), and failure of a  $\geq$  3 consecutive month trial of at least ONE conventional DMARD (e.g., sulfasalazine, leflunomide, hydroxychloroquine) at up to maximally indicated doses, unless clinically significant adverse effects are experienced or all are contraindicated;

## CLINICAL POLICY

### Anakinra

5. Failure of ALL of the following, each used for  $\geq 3$  consecutive months, unless clinically significant adverse effects are experienced or all are contraindicated (a and b):
  - a. Enbrel<sup>®</sup>;
  - b. Member has not responded or is intolerant to one or more TNF blockers (i.e. Xeljanz<sup>®</sup>) unless member has cardiovascular risk and benefits do not outweigh the risk of treatment;  
*\*Prior authorization may be required for Enbrel and Xeljanz*
6. Documentation of one of the following baseline assessment scores (a or b):
  - a. Clinical disease activity index (CDAI) score (*see Appendix G*);
  - b. Routine assessment of patient index data 3 (RAPID3) score (*see Appendix H*);
7. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
8. Dose does not exceed 100 mg per day.

**Approval duration: 6 months**

#### **B. Cryopyrin-Associated Periodic Syndromes (must meet all):**

1. Diagnosis of NOMID or chronic infantile neurological, cutaneous and articular syndrome (CINCA);
2. Prescribed by or in consultation with a rheumatologist;
3. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
4. Dose does not exceed 8 mg/kg per day (*see Appendix E for dose rounding guidelines*).

**Approval duration: 6 months**

#### **C. Deficiency of Interleukin-1 Receptor Antagonist (must meet all):**

1. Diagnosis of DIRA confirmed by presence of loss-of-function *ILRN* mutations;
2. Prescribed by or in consultation with a rheumatologist;
3. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
4. Dose does not exceed 8 mg/kg per day (*see Appendix E for dose rounding guidelines*).

**Approval duration: 6 months**

#### **D. Other diagnoses/indications**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

## **II. Continued Therapy**

### **A. All Indications in Section I (must meet all):**

## CLINICAL POLICY

### Anakinra

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member meets one of the following (a or b):
  - a. For RA: member is responding positively to therapy as evidenced by one of the following (i or ii):
    - i. A decrease in CDAI (*see Appendix G*) or RAPID3 (*see Appendix H*) score from baseline;
    - ii. Medical justification stating inability to conduct CDAI re-assessment, and submission of RAPID3 score associated with disease severity that is similar to initial CDAI assessment or improved;
  - b. For all other indications: member is responding positively to therapy;
3. Member does not have combination use with biological disease-modifying antirheumatic drugs or Janus kinase inhibitors (*see Section III: Diagnoses/Indications for which coverage is NOT authorized*);
4. If request is for a dose increase, new dose does not exceed one of the following (a or b):
  - a. RA: 100 mg per day;
  - b. DIRA, NOMID: 8 mg/kg per day (*see Appendix E for dose rounding guidelines*).

**Approval duration: 12 months**

#### **B. Other diagnoses/indications (must meet 1 or 2):**

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.  
**Approval duration: Duration of request or 6 months (whichever is less);** or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

### **III. Diagnoses/Indications for which coverage is NOT authorized:**

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.PMN.53 for Medicaid or evidence of coverage documents;
- B.** Combination use with biological disease-modifying antirheumatic drugs (bDMARDs) or potent immunosuppressants, including but not limited to any tumor necrosis factor (TNF) antagonists [e.g., Cimzia<sup>®</sup>, Enbrel<sup>®</sup>, Humira<sup>®</sup>, Simponi<sup>®</sup>, Avsola<sup>™</sup>, Inflectra<sup>™</sup>, Remicade<sup>®</sup>, Renflexis<sup>™</sup>], interleukin agents [e.g., Arcalyst<sup>®</sup> (IL-1 blocker), Ilaris<sup>®</sup> (IL-1 blocker), Kineret<sup>®</sup> (IL-1RA), Actemra<sup>®</sup> (IL-6RA), Kevzara<sup>®</sup> (IL-6RA), Stelara<sup>®</sup> (IL-12/23 inhibitor), Cosentyx<sup>®</sup> (IL-17A inhibitor), Taltz<sup>®</sup> (IL-17A inhibitor), Siliq<sup>™</sup> (IL-17RA), Ilumya<sup>™</sup> (IL-23 inhibitor), Skyrizi<sup>™</sup> (IL-23 inhibitor), Tremfya<sup>®</sup> (IL-23 inhibitor)], Janus kinase inhibitors (JAKi) [e.g., Xeljanz<sup>®</sup>/Xeljanz<sup>®</sup> XR, Cibinqo<sup>™</sup>, Olumiant<sup>™</sup>, Rinvoq<sup>™</sup>], anti-CD20 monoclonal antibodies [Rituxan<sup>®</sup>, Riabni<sup>™</sup>, Ruxience<sup>™</sup>, Truxima<sup>®</sup>, Rituxan Hycela<sup>®</sup>], selective co-stimulation modulators [Orencia<sup>®</sup>], and integrin receptor antagonists [Entyvio<sup>®</sup>] because of the additive immunosuppression, increased risk of neutropenia, as well as increased risk of serious infections.

## CLINICAL POLICY

### Anakinra

#### IV. Appendices/General Information

##### Appendix A: Abbreviation/Acronym Key

CAPS: cryopyrin-associated periodic syndromes

CDAI: clinical disease activity index

DIRA: deficiency of Interleukin-1

Receptor Antagonist

DMARD: disease-modifying

antirheumatic drug

FDA: Food and Drug Administration

IL-1: interleukin-1

JAKi: Janus kinase inhibitors

MTX: methotrexate

NOMID: neonatal-onset multisystem inflammatory disease

RA: rheumatoid arthritis

RAPID3: routine assessment of patient index data 3

##### Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
azathioprine (Azasan <sup>®</sup> , Imuran <sup>®</sup> )	<b>RA</b> 1 mg/kg/day PO QD or divided BID	2.5 mg/kg/day
Cuprimine <sup>®</sup> (d-penicillamine)	<b>RA*</b> <u>Initial dose:</u> 125 or 250 mg PO QD <u>Maintenance dose:</u> 500 – 750 mg/day PO QD	1,500 mg/day
cyclosporine (Sandimmune <sup>®</sup> , Neoral <sup>®</sup> )	<b>RA</b> 2.5 – 4 mg/kg/day PO divided BID	4 mg/kg/day
hydroxychloroquine (Plaquenil <sup>®</sup> )	<b>RA*</b> <u>Initial dose:</u> 400 – 600 mg/day PO QD <u>Maintenance dose:</u> 200 – 400 mg/day PO QD	600 mg/day
leflunomide (Arava <sup>®</sup> )	<b>RA</b> 100 mg PO QD for 3 days, then 20 mg PO QD	20 mg/day
methotrexate (Rheumatrex <sup>®</sup> )	<b>RA</b> 7.5 mg/week PO, SC, or IM or 2.5 mg PO Q12 hr for 3 doses/week	30 mg/week
Ridaura <sup>®</sup> (auranofin)	<b>RA</b> 6 mg PO QD or 3 mg PO BID	9 mg/day (3 mg TID)
sulfasalazine (Azulfidine <sup>®</sup> )	<b>RA</b> 2 g/day PO in divided doses	3 g/day
Actemra <sup>®</sup> (tocilizumab)	<b>RA</b>	IV: 800 mg every 4 weeks

## CLINICAL POLICY

### Anakinra

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	IV: 4 mg/kg every 4 weeks followed by an increase to 8 mg/kg every 4 weeks based on clinical response  SC: Weight < 100 kg: 162 mg SC every other week, followed by an increase to every week based on clinical response Weight ≥ 100 kg: 162 mg SC every week	SC: 162 mg every week
Enbrel® (etanercept)	<b>RA</b> 25 mg SC twice weekly or 50 mg SC once weekly	50 mg/week
Kevzara® (sarilumab)	<b>RA</b> 200 mg SC once every two weeks	200 mg/2 weeks
Oluminat® (baricitinib)	<b>RA</b> 2 mg PO QD	2 mg/day
Xeljanz® (tofacitinib)	<b>RA</b> 5 mg PO BID	10 mg/day
Xeljanz XR® (tofacitinib extended-release)	<b>RA</b> 11 mg PO QD	11 mg/day

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

\*Off-label

#### Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): known hypersensitivity to *E. coli*-derived proteins, Kineret, or any components of the product
- Boxed warning(s): none reported

#### Appendix D: General Information

- Definition of MTX or DMARD Failure
  - Child-bearing age is not considered a contraindication for use of MTX. Each drug has risks in pregnancy. An educated patient and family planning would allow use of MTX in patients who have no intention of immediate pregnancy.
  - Social use of alcohol is not considered a contraindication for use of MTX. MTX may only be contraindicated if patients choose to drink over 14 units of alcohol per week. However, excessive alcohol drinking can lead to worsening of the condition, so patients who are serious about clinical response to therapy should refrain from excessive alcohol consumption.
- Examples of positive response to therapy may include, but are not limited to:
  - Reduction in joint pain/swelling/tenderness
  - Improvement in ESR/CRP levels

## CLINICAL POLICY

### Anakinra

- Improvements in activities of daily living
- DIRA patients are homozygous or compound heterozygous for loss-of-function mutations in *IL1RN*, encoding IL-1Ra. Most mutations are nonsense or frameshift mutations that lead to either no expression of protein or expression of nonfunctional protein. Examples of disease-causing mutations in *IL1RN* identified include: 4 nonsense mutations, 1 in-frame deletion, 3 frameshift deletions, and a 22-kb and a genomic 175-kb deletion on chromosome 2.

#### Appendix E: Dose Rounding Guidelines for DIRA, NOMID

Weight-based Dose Range	Vial Quantity Recommendation
≤ 104.99 mg	1 syringe of 100 mg/0.67 mL
105 to 209.99 mg	2 syringes of 100 mg/0.67 mL
210 to 314.99 mg	3 syringes of 100 mg/0.67 mL
315 to 419.99 mg	4 syringes of 100 mg/0.67 mL
420 to 524.99 mg	5 syringes of 100 mg/0.67 mL
525 to 629.99 mg	6 syringes of 100 mg/0.67 mL
630 to 734.99 mg	7 syringes of 100 mg/0.67 mL
735 to 839.99 mg	8 syringes of 100 mg/0.67 mL

#### Appendix F: The 2010 ACR Classification Criteria for RA

Add score of categories A through D; a score of ≥ 6 out of 10 is needed for classification of a patient as having definite RA.

A	Joint involvement	Score
	1 large joint	0
	2-10 large joints	1
	1-3 small joints (with or without involvement of large joints)	2
	4-10 small joints (with or without involvement of large joints)	3
	> 10 joints (at least one small joint)	5
B	Serology (at least one test result is needed for classification)	
	Negative rheumatoid factor (RF) and negative anti-citrullinated protein antibody (ACPA)	0
	Low positive RF or low positive ACPA * Low: < 3 x upper limit of normal	2
	High positive RF or high positive ACPA * High: ≥ 3 x upper limit of normal	3
C	Acute phase reactants (at least one test result is needed for classification)	
	Normal C-reactive protein (CRP) and normal erythrocyte sedimentation rate (ESR)	0
	Abnormal CRP or abnormal ESR	1
D	Duration of symptoms	
	< 6 weeks	0
	≥ 6 weeks	1

#### Appendix G: Clinical Disease Activity Index (CDAI) Score

## CLINICAL POLICY

### Anakinra

The Clinical Disease Activity Index (CDAI) is a composite index for assessing disease activity in RA. CDAI is based on the simple summation of the count of swollen/tender joint count of 28 joints along with patient and physician global assessment on VAS (0–10 cm) Scale for estimating disease activity. The CDAI score ranges from 0 to 76.

CDAI Score	Disease state interpretation
≤ 2.8	Remission
> 2.8 to ≤ 10	Low disease activity
> 10 to ≤ 22	Moderate disease activity
> 22	High disease activity

#### Appendix H: Routine Assessment of Patient Index Data 3 (RAPID3) Score

The Routine Assessment of Patient Index Data 3 (RAPID3) is a pooled index of the three patient-reported ACR core data set measures: function, pain, and patient global estimate of status. Each of the individual measures is scored 0 – 10, and the maximum achievable score is 30.

RAPID3 Score	Disease state interpretation
≤ 3	Remission
3.1 to 6	Low disease activity
6.1 to 12	Moderate disease activity
> 12	High disease activity

## V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
RA	100 mg SC QD	100 mg/day
NOMID	<u>Initial dose:</u> 1 – 2 mg/kg SC QD or divided BID <u>Maintenance dose:</u> Adjust doses in 0.5 to 1 mg/kg increments. Once daily administration is recommended, but the dose may be split into twice daily administration (a new syringe must be used for each dose).	8 mg/kg/day
DIRA	<u>Initial dose:</u> 1 – 2 mg/kg SC QD <u>Maintenance dose:</u> Adjust doses in 0.5 to 1 mg/kg increments.	8 mg/kg/day

## VI. Product Availability

Single-use prefilled syringe: 100 mg/0.67 mL

## VII. References

1. Kineret Prescribing Information. Stockholm, Sweden: Swedish Orphan Biovitrum AB; December 2020. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/103950s51891bl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/103950s51891bl.pdf). Accessed February 20, 2022.



## CLINICAL POLICY

### Anakinra

2. Fraenkel L, Bathon JM, Enggland BR, et al. 2021 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care & Research*. 2021; 73(7):924-939. DOI 10.1002/acr.24596.
3. Chiabrando JG, Bonaventura A, Vecchie A, et al. Management of acute and recurrent pericarditis: JACC State-of-the-Art Review. *J Am Coll Cardiol* 2020; 75(1):76-92. <http://doi.org/10.1016/j.jacc.2019.11.021>.

### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J3590	Unclassified biologics

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	01.23	01.23

### Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or



## CLINICAL POLICY

### Anakinra

regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.