

## Clinical Policy: Calcipotriene/Betamethasone Dipropionate Foam (Enstilar)

Reference Number: DE.PMN.181

Effective Date: 01.23

Last Review Date: 01.23

Line of Business: Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

Calcipotriene 0.005% and betamethasone dipropionate 0.064% foam (Enstilar<sup>®</sup>) is a combination topical product of a vitamin D analog and a corticosteroid.

### FDA Approved Indication(s)

Enstilar is indicated for the topical treatment of plaque psoriasis (PsO) in patients 12 years of age and older.

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Enstilar is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Plaque Psoriasis (must meet all):

1. Diagnosis of PsO;
2. Age  $\geq$  12 years;
3. Failure of a medium to ultra-high potency topical corticosteroid (*see Appendix B*) unless contraindicated or clinically significant adverse effects are experienced;
4. Failure of the following, unless clinically significant adverse effects are experienced or contraindicated: calcipotriene;
5. Dose does not exceed 60 g every 4 days (7 canisters per month).

**Approval duration: One month**

##### B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

#### II. Continued Therapy

##### A. Plaque Psoriasis (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy;

## CLINICAL POLICY

### Calcipotriene/Betamethasone Dipropionate Foam

3. If request is for a dose increase, new dose does not exceed 60 gm every 4 days (7 canisters per month).

**Approval duration: Up to one month of total treatment** (a single continuous course of therapy up to 4 weeks is recommended)

#### B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

**Approval duration: Duration of request or 1 month (whichever is less);** or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

### III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.PMN.53 for Medicaid or evidence of coverage documents.

### IV. Appendices/General Information

*Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

PsO: plaque psoriasis

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
calcipotriene (Dovonex <sup>®</sup> ) cream, ointment, solution	Apply topically to the affected area(s) BID	100 g/week
calcitriol (Vectical <sup>™</sup> ) ointment	Apply topically to the affected area(s) BID	200 g/week
tazarotene (Tazorac <sup>®</sup> ) gel, cream	Apply topically to the affected area(s) QHS	Once daily application
<b>Ultra-High Potency Topical Corticosteroids</b>		
augmented betamethasone dipropionate 0.05% (Diprolene <sup>®</sup> , Alphatrex <sup>®</sup> ) ointment, gel	Apply topically to the affected area(s) BID	Should not be used for longer than 2 consecutive weeks
clobetasol propionate 0.05% (Temovate <sup>®</sup> , Temovate E <sup>®</sup> ) cream, ointment, gel, solution		
diflorasone diacetate 0.05% (Apexicon <sup>®</sup> ) ointment		
halobetasol propionate 0.05% (Ultravate <sup>®</sup> ) cream, ointment		
<b>High Potency Topical Corticosteroids</b>		

## CLINICAL POLICY

### Calcipotriene/Betamethasone Dipropionate Foam

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
augmented betamethasone dipropionate 0.05% (Diprolone <sup>®</sup> , Diprolene <sup>®</sup> AF) cream, lotion	Apply topically to the affected area(s) BID	Should not be used for longer than 2 consecutive weeks
betamethasone dipropionate 0.05% ointment		
desoximetasone (Topicort <sup>®</sup> ) 0.25%, 0.05% cream, ointment, gel		
diflorasone 0.05% (Apexicon E <sup>®</sup> ) cream		
fluocinonide acetone 0.05% cream, ointment, gel, solution		
triamcinolone acetone 0.5% (Aristocort <sup>®</sup> , Kenalog <sup>®</sup> ) cream, ointment		
Medium/Medium to High Potency Topical Corticosteroids		
betamethasone dipropionate 0.05% cream	Apply topically to the affected area(s) BID	Should not be used for longer than 2 consecutive weeks
desoximetasone 0.05% (Topicort <sup>®</sup> ) cream, ointment, gel		
fluocinolone acetone 0.025% (Synalar <sup>®</sup> ) cream, ointment		
fluticasone propionate 0.05% (Cutivate <sup>®</sup> ) cream		
mometasone furoate 0.1% (Elocon <sup>®</sup> ) cream, lotion, ointment		
triamcinolone acetone 0.1%, 0.25%, 0.5% (Aristocort <sup>®</sup> , Kenalog <sup>®</sup> ) cream, ointment		

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

#### Appendix C: Contraindications/Boxed Warnings

None reported

## V. Dosage and Administration

Drug Name	Dosing Regimen	Maximum Dose
calcipotriene 0.005%/betamethasone dipropionate 0.064% (Enstilar)	Apply topically to affected area(s) QD for up to 4 weeks. Avoid use on face, groin, axillae, skin treatment site with atrophy present, or with occlusive dressing unless directed by a healthcare provider.	60 g/4 days

## VI. Product Availability

Can containing 0.005% calcipotriene/0.064% betamethasone dipropionate foam: 60 g, 120 g (2 cans of 60 g)

## VII. References

1. Enstilar Prescribing Information. Parsippany, NJ: LEO Laboratories Ltd; April 2022. Available at: <https://www.enstilar.com>. Accessed July 20, 2022.

## CLINICAL POLICY

### Calcipotriene/Betamethasone Dipropionate Foam

2. Elmets CA, Korman NJ, Prater EF, et al. Joint AAD-NPF Guidelines of care for the management and treatment of psoriasis with topical therapy and alternative medicine modalities for psoriasis severity measures. J Am Acad Dermatol 2021;84:432-70.
3. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2022. Available at: <http://www.clinicalpharmacology-ip.com/>. Accessed July 20, 2022.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	01.23	01.23

#### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

## CLINICAL POLICY

### Calcipotriene/Betamethasone Dipropionate Foam

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.