

## Clinical Policy: Sacituzumab Govitecan-hziy (Trodelvy)

Reference Number: CP.PHAR.475

Effective Date: 04.22.20

Last Review Date: 05.22

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)  
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

Sacituzumab govitecan-hziy (Trodelvy<sup>™</sup>) is a Trop-2-directed antibody and topoisomerase inhibitor conjugate.

### FDA Approved Indication(s)

Trodelvy is indicated for the treatment of adult patients with

- Unresectable locally advanced or metastatic triple-negative breast cancer (mTNBC) who have received two or more prior systemic therapies, at least one of them for metastatic disease
- Locally advanced or metastatic urothelial cancer who have previously received a platinum-containing chemotherapy and either programmed death receptor-1 (PD-1) or programmed death-ligand 1 (PDL1) inhibitor\*

---

*\*This indication is approved under accelerated approval based on tumor response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.*

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Trodelvy is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Breast Cancer (must meet all):

1. Diagnosis of unresectable or metastatic breast cancer;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq$  18 years;
4. Documentation of triple negative (i.e., estrogen receptor-, progesterone receptor-, and human epidermal growth factor receptor 2 [HER2]-negative) disease;
5. Failure of both of the following (a and b):
  - a. Two or more prior regimens (*see Appendix B*);
  - b. At least one of the prior regimens administered for metastatic disease (*see Appendix B*);
6. Prescribed as a single agent;
7. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 10 mg/kg on days 1 and 8 of each 21-day cycle;

- b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration:**

**HIM/Medicaid** – 6 months

**Commercial** – 6 months or to the member’s renewal date

**B. Urothelial Cancer (must meet all):**

1. Diagnosis of locally advanced, recurrent, or metastatic urothelial cancer;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq$  18 years;
4. Failure of both of the following (a and b):
  - a. Platinum-containing chemotherapy (*see Appendix B*);
  - b. Programmed death receptor-1 (PD-1) or programmed death-ligand 1 (PD-L1) inhibitor (*see Appendix B*);
5. Prescribed as a single agent;
6. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 10 mg/kg on days 1 and 8 of each 21-day cycle;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration:**

**HIM/Medicaid** – 6 months

**Commercial** – 6 months or to the member’s renewal date

**C. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

## II. Continued Therapy

### A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Trodelvy for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):\*
  - a. New dose does not exceed 10 mg/kg on days 1 and 8 of each 21-day cycle;
  - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

\*Prescribed regimen must be FDA-approved or recommended by NCCN

### Approval duration:

**HIM/Medicaid** – 12 months

**Commercial** – 6 months or to the member's renewal date

### B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

## III. Diagnoses/Indications for which coverage is NOT authorized:

- ### A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

## IV. Appendices/General Information

### Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

HER2: human epidermal growth factor receptor 2

PD-1: programmed death receptor-1

PD-L1: programmed death-ligand

TNBC: triple-negative breast cancer

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
<b>Examples of systemic therapies for recurrent unresectable or metastatic breast cancer</b>		
paclitaxel	Varies	Varies
Abraxane <sup>®</sup> (albumin-bound paclitaxel)	Varies	Varies
docetaxel (Taxotere <sup>®</sup> )	Varies	Varies
doxorubicin	Varies	Varies
Liposomal doxorubicin (Doxil <sup>®</sup> )	50 mg/m <sup>2</sup> IV day 1, cycled every 28 days	Varies
capecitabine (Xeloda <sup>®</sup> )	1,000-1,250 mg/m <sup>2</sup> PO BID on days 1-14, cycled every 21 days	Varies
gemcitabine (Gemzar <sup>®</sup> )	800-1,200 mg/m <sup>2</sup> IV on days 1,8 and 15, cycled every 28 days	Varies
vinorelbine	Varies	Varies
Halaven <sup>®</sup> (eribulin)	1.4 mg/m <sup>2</sup> IV on days 1 and 8, cycled every 21 days	Varies
carboplatin	AUC 6 IV on day 1, cycled every 21-28 days	Varies
cisplatin	75 mg/m <sup>2</sup> IV on day 1, cycled every 21 days	Varies
cyclophosphamide	50 mg PO QD on days 1-21, cycled every 28 days	Varies
epirubicin (Ellence <sup>®</sup> )	60-90 mg/m <sup>2</sup> IV on day 1, cycled every 21 days	Varies
Ixempra <sup>®</sup> (ixabepilone)	40 mg/m <sup>2</sup> IV on day 1, cycled every 21 days	40 mg/m <sup>2</sup>
<b>Examples of platinum-containing regimens for urothelial cancer</b>		
DDMVAC (dose-dense methotrexate, vinblastine, doxorubicin, and cisplatin)	Varies	Varies
gemcitabine with either cisplatin or carboplatin	Varies	Varies
<b>Examples of PD-1 and PD-L1 inhibitors for urothelial cancer</b>		
Keytruda <sup>®</sup> (pembrolizumab)	Varies	Varies
Tecentriq <sup>®</sup> (atezolizumab)	Varies	Varies
Opdivo <sup>®</sup> (nivolumab)	Varies	Varies
Bavencio <sup>®</sup> (avelumab)	800 mg IV infusion once every 2 weeks	Varies

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): severe hypersensitivity reaction to Trodelvy
- Boxed warning(s): neutropenia and diarrhea

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
Triple- negative breast cancer, urothelial cancer	10 mg/kg IV on days 1 and 8 of each 21-day cycle	10 mg/kg

**VI. Product Availability**

Single-dose vial: 180 mg lyophilized powder for reconstitution

**VII. References**

1. Trodelvy Prescribing Information. Morris Plains, NJ: Immunomedics, Inc.; October 2021. Available at: [https://www.gilead.com/-/media/files/pdfs/medicines/oncology/trodelvy/trodelvy\\_pi.pdf](https://www.gilead.com/-/media/files/pdfs/medicines/oncology/trodelvy/trodelvy_pi.pdf). Accessed February 13, 2022.
2. Bardia A, Mayer IA, Vahdat LT, et al. Sacituzumab Govitecan-hziy in refractory metastatic triple-negative breast cancer. *N Engl J Med* 2019 Feb 21;380(8):741-51.
3. Sacituzumab govitecan. In: National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: [http://www.nccn.org/professionals/drug\\_compendium](http://www.nccn.org/professionals/drug_compendium). Accessed February 13, 2022.

**Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9317	Injection, sacituzumab govitecan-hziy, 2.5 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created pre-emptively.	03.03.20	05.20
Drug is now FDA-approved - criteria updated per FDA-labeling: removed requirement for previous taxane-based regimen as this is neither in the PI nor required by NCCN.	05.10.20	08.20
2Q 2021 annual review: RT4: added criteria for new mUC indication; updated breast cancer criteria to add unresectable locally advanced option and clarified that of the two or more prior regimens, at least one of them be for metastatic disease, based on updated FDA-labeling; updated JCode; references for HIM line of business off-label use revised from HIM.PHAR.21 to HIM.PA.154; references reviewed and updated.	04.19.21	05.21
2Q 2022 annual review: for TNBC: removed “locally advanced” requirement as disease can be local or regional per NCCN; added recurrent urothelial carcinoma indication per NCCN; added criterion	02.13.22	05.22

Reviews, Revisions, and Approvals	Date	P&T Approval Date
for use as single-agent therapy for both TNBC and urothelial cancer per NCCN; references reviewed and updated.		
Template changes applied to other diagnoses/indications.	09.28.22	

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2020 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.