

Clinical Policy: Tralokinumab-ldrm (Adbry)

Reference Number: DE.PHAR.577

Effective Date: 01.23

Last Review Date: 01.23

Line of Business: Medicaid

[Coding Implications](#)
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Tralokinumab-ldrm (Adbry[®]) is an interleukin-13 antagonist.

FDA Approved Indication(s)

Adbry is indicated for the treatment of moderate-to-severe atopic dermatitis in adult patients whose disease is not adequately controlled with topical prescription therapies or when those therapies are not advisable. Adbry can be used with or without topical corticosteroids.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Adbry is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Atopic Dermatitis (must meet all):

1. Diagnosis of atopic dermatitis affecting one of the following (a or b):
 - a. At least 10% of the member's body surface area (BSA);
 - b. Hands, feet, face, neck, scalp, genitals/groin, and/or intertriginous areas;
2. Prescribed by or in consultation with a dermatologist or allergist;
3. Age \geq 18 years;
4. Failure of all of the following (a, b, and c), unless contraindicated or clinically significant adverse effects are experienced:
 - a. One formulary medium to very high potency topical corticosteroids, each used for \geq 2 weeks;
 - b. One non-steroidal topical therapy* used for \geq 4 weeks: topical calcineurin inhibitor (e.g., tacrolimus 0.03% ointment, pimecrolimus 1% cream) or Eucrisa[®];
**These agents may require prior authorization*
 - c. One systemic agent used for \geq 3 months: azathioprine, methotrexate, mycophenolate mofetil, or cyclosporine;
5. Adbry is not prescribed concurrently with another biologic medication (e.g., Dupixent[®]) or a JAK inhibitor (e.g., Olumiant[®], Rinvoq[®], Cibinqo[®], Opzelura[™]);
6. Dose does not exceed the following:
 - a. Initial (one-time) dose of 600 mg (four injections);
 - b. Maintenance dose of 300 mg (two injections) every 2 weeks;

CLINICAL POLICY

Tralokinumab-ldrm

Approval duration: 4 months (18 injections)

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53

II. Continued Therapy

A. Atopic Dermatitis (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by, including but not limited to, reduction in itching and scratching;
3. Adbry is not prescribed concurrently with another biologic medication (e.g., Dupixent) or a JAK inhibitor (e.g., Olumiant, Rinvoq, Cibinqo, Opzelura);
4. For members with weight < 100 kg: Request is for 300 mg every 4 weeks, unless documentation supports member has not achieved clear or almost clear skin;
5. If request is for a dose increase, new dose does not exceed 300 mg every 2 weeks.

Approval duration:

Medicaid – 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 6 months (whichever is less); or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53

III. Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – CP.PMN.53 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

BSA: body surface area

FDA: Food and Drug Administration

JAK: Janus kinase

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

CLINICAL POLICY
Tralokinumab-ldrm

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Very High Potency Topical Corticosteroids		
augmented betamethasone 0.05% (Diprolene® AF) cream, ointment, gel, lotion	Apply topically to the affected area(s) BID	Varies
clobetasol propionate 0.05% (Temovate®) cream, ointment, gel, solution		
diflorasone diacetate 0.05% (Maxiflor®, Psorcon E®) cream, ointment		
halobetasol propionate 0.05% (Ultravate®) cream, ointment		
High Potency Topical Corticosteroids		
augmented betamethasone 0.05% (Diprolene® AF) cream, ointment, gel, lotion	Apply topically to the affected area(s) BID	Varies
diflorasone 0.05% (Florone®, Florone E®, Maxiflor®, Psorcon E®) cream		
fluocinonide acetone 0.05% (Lidex®, Lidex E®) cream, ointment, gel, solution		
triamcinolone acetonide 0.5% (Aristocort®, Kenalog®) cream, ointment		
Medium Potency Topical Corticosteroids		
desoximetasone 0.05% (Topicort®) cream, ointment, gel	Apply topically to the affected area(s) BID	Varies
fluocinolone acetonide 0.025% (Synalar®) cream, ointment		
mometasone 0.1% (Elocon®) cream, ointment, lotion		
triamcinolone acetonide 0.025%, 0.1% (Aristocort®, Kenalog®) cream, ointment		
Low Potency Topical Corticosteroids		
alclometasone 0.05% (Aclovate®) cream, ointment	Apply topically to the affected area(s) BID	Varies
desonide 0.05% (Desowen®) cream, ointment, lotion		
fluocinolone acetonide 0.01% (Synalar®) solution		
hydrocortisone 2.5% (Hytone®) cream, ointment		
Other Classes of Agents		
Protopic® (tacrolimus), Elidel® (pimecrolimus)	Children ≥ 2 years and adults: Apply a thin	Varies

CLINICAL POLICY

Tralokinumab-ldrm

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	layer topically to affected skin BID. Treatment should be discontinued if resolution of disease occurs.	
Eucrisa® (crisaborole)	Apply to the affected areas BID	Varies
cyclosporine	3-6 mg/kg/day PO BID	300 mg/day
azathioprine	1-3 mg/kg/day PO QD	Weight-based
methotrexate	7.5-25 mg/wk PO once weekly	25 mg/week
mycophenolate mofetil	1-1.5 g PO BID	3 g/day

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): known hypersensitivity to tralokinumab-ldrm or any excipients in Adbry
- Boxed warning(s): none

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Moderate-to-severe atopic dermatitis	Initial dose of 600 mg SC followed by 300 mg SC every other week After 16 weeks of treatment, for patients with body weight < 100 kg who achieve clear or almost clear skin, a dosage of 300 mg every 4 weeks may be considered	See regimen

VI. Product Availability

Pre-filled syringe: 150 mg/mL

VII. References

1. Adbry Prescribing Information. Madison, NJ: LEO Pharma, Inc.; December 2021. Available at: <https://www.adbry.com/>. Accessed January 20, 2022.
2. DRUGDEX® System [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed January 20, 2022.
3. Wollenberg A, Christen-Zäch S, Taieb A, et al. ETFAD/EADV Eczema task force 2020 position paper on diagnosis and treatment of atopic dermatitis in adults and children. *J Eur Acad Dermatol Venereol*. 2020 Dec;34(12):2717-2744.
4. Eichenfield F, Tom WL, Chamlin SL, et al. Guidelines of Care for the Management of Atopic Dematitis. *J Am Acad Dermatol*. 2014 February; 70(2): 338–351.

CLINICAL POLICY

Tralokinumab-ldrm

5. Wollenberg A, Blauvelt A, Guttman-Yassky E, et al. Tralokinumab for moderate-to-severe atopic dermatitis: results from two 52-week, randomized, double-blind, multicentre, placebo-controlled phase III trials (ECZTRA 1 and ECZTRA 2). *Br J Dermatol.* 2021 Mar;184(3):437-449.
6. Silverberg JI, Toth D, Bieber T, et al. Tralokinumab plus topical corticosteroids for the treatment of moderate-to-severe atopic dermatitis: results from the double-blind, randomized, multicentre, placebo-controlled phase III ECZTRA 3 trial. *Br J Dermatol.* 2021 Mar;184(3):450-463.
7. Drucker AM, Ellis AG, Bohdanowicz M, et al. Systemic Immunomodulatory Treatments for Patients with Atopic Dermatitis: A Systematic Review and Network Meta-analysis. *JAMA Dermatol.* 2020;156(6):659-667.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J3590	Unclassified biologics
C9399	Unclassified drugs or biologicals (hospital outpatient use)

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	09.22	11.22

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy,

CLINICAL POLICY

Tralokinumab-ldrm

contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2022 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.